

Author's Accepted Manuscript

A SYSTEMATIC REVIEW AND META-ANALYSIS OF RANDOMIZED AND NON-RANDOMIZED TRIALS OF CLINICAL EMOTIONAL FREEDOM TECHNIQUES (EFT) FOR THE TREATMENT OF DEPRESSION

Jerrold A. Nelms, Liana Castel



PII: S1550-8307(16)30106-9
DOI: <http://dx.doi.org/10.1016/j.explore.2016.08.001>
Reference: JSCH2129

To appear in: *Explore: The Journal of Science and Healing*

Received date: 2 October 2015

Cite this article as: Jerrod A. Nelms and Liana Castel, A SYSTEMATIC REVIEW AND META-ANALYSIS OF RANDOMIZED AND NON-RANDOMIZED TRIALS OF CLINICAL EMOTIONAL FREEDOM TECHNIQUES (EFT) FOR THE TREATMENT OF DEPRESSION, *Explore: The Journal of Science and Healing*, <http://dx.doi.org/10.1016/j.explore.2016.08.001>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

A Systematic Review and Meta-Analysis of Randomized and Non-Randomized Trials of Clinical Emotional Freedom Techniques (EFT) for the Treatment of Depression

Citation: Nelms, J., & Castel D. (2015). A Systematic Review and Meta-Analysis of Randomized and Non-Randomized Trials of Emotional Freedom Techniques (EFT) for the Treatment of Depression [insert journal, volume, number, and pages here]

Jerrod A. Nelms, Ph.D., MPH¹

Liana Castel, Ph.D., MSPH¹

1. Lucyna Health and Safety Solutions, LLC, Tampa, FL

Corresponding author: Jerrod Nelms

Email: jnelms@lucynasolutions.com

Phone: +1-813-335-9640

Abstract

Background: Among a group of therapies collectively known as Energy Psychology (EP), Emotional Freedom Techniques (EFT) is the most widely practiced. Clinical EFT is an evidence-based practice combining elements of cognitive and exposure therapies with the manual stimulation of acupuncture points (acupoints). Lacking is a recent quantitative meta-analysis that enhances understanding of the variability and clinical significance of depression reduction outcomes after Clinical EFT treatment.

Methods: All studies (2005–2015) evaluating EFT for sufferers of depression were identified by electronic search; these included both outcome studies and randomized controlled trials (RCTs). Our focus was depressive symptoms as measured by a variety of psychometric questionnaires

and scales. We used meta-analysis to calculate effect sizes at three time points including posttest, follow-ups less than 90 days, and follow-ups greater than 90 days.

Results: Twenty studies qualified for inclusion, 12 RCTs and 8 outcome studies. The number of participants treated with EFT included $N = 461$ in outcome studies and $N = 398$ in RCTs.

Clinical EFT showed a large effect size in the treatment of depression in RCTs. At posttest Cohen's d for RCTs was 1.85 and for outcome studies was 0.70. Effect sizes for follow-ups less than 90 days was 1.21, and for ≥ 90 days was 1.11. EFT was more efficacious than DB (Diaphragmatic Breathing) and SI (Supportive Interview) in posttest measurements ($p = 0.06$ vs DB; $p < 0.001$ vs SI), and SHE (Sleep Hygiene Education) at follow-up ($p = 0.036$). No significant treatment effect difference between EFT and EMDR (Eye Movement Desensitization and Reprocessing) was found. EFT was superior to TAU (treatment as usual), and efficacious in treatment time frames ranging from one to 10 sessions. The mean of symptom reductions across all studies was -41%.

Conclusion: The results show that Clinical EFT is highly effective in reducing depressive symptoms in a variety of populations and settings. EFT was equal or superior to TAU and other active treatment controls. The posttest effect size for EFT ($d = 1.31$) was larger than that measured in meta-analyses of antidepressant drug trials and psychotherapy studies. EFT produced large treatment effects whether delivered in group or individual format, and participants maintained their gains over time. This meta-analysis extends the existing literature through facilitation of a better understanding of the variability and clinical significance of depression improvement subsequent to EFT treatment.

Background

Emotional Freedom Techniques (EFT) is an evidence-based practice, with more than 100 randomized controlled trials (RCTs), outcome studies, and review papers published in peer-reviewed journals listed in an online research bibliography (Research.EFTuniverse.com). EFT meets the criteria of the American Psychological Association's Division 12 Task Force on Empirically Validated Treatments (hereafter referred to as APA Division 12 criteria) for a number of psychological conditions, including anxiety, depression, phobias, and PTSD (Church, Feinstein, Palmer-Hoffman, Stein, & Tranguch, 2014). EFT has also been found to produce improvement in physical conditions such as fibromyalgia (Brattberg, 2008), psoriasis (Hodge & Jurgens, 2011), tension headaches (Bougea et al., 2013), pain (Church, 2014), traumatic brain injury (Church & Brooks, 2014), and seizure disorders (Swingle, 2010). In service evaluations performed by the UK National Health Service, EFT has been found to produce an improvement in general physical functioning as well as mental health (Stewart, et al., 2013; Boath, Carryer & Stewart, 2013; Stewart, Boath, Carryer, Walton, & Hill, 2014). The wide range of conditions for which EFT is effective is usually attributed to the technique's ability to reduce stress, which is a component of many emotional and physical disorders (Lane, 2009; Church, 2013a).

EFT treatment is standardized via a manual, with most published studies applying the manualized form of the technique (Craig & Fowlie, 1995; Church, 2013b). EFT combines acupressure with elements of two other evidence-based psychotherapeutic techniques, namely exposure therapy and cognitive therapy. While vividly recalling a traumatic event, clients are instructed to tap on 12 acupressure points with their fingertips. The client reports the emotional intensity of the event on an 11-point Likert scale (after Wolpe, 1958) both before and after the

application of EFT. Studies and clinical reports note rapid reductions in distress after tapping, client safety, and an absence of adverse events after treatment (Church, 2013a; Mollon, 2007; Schulz, 2009). Clinical EFT is the manualized evidence-based form of the original EFT methodology (Church, 2013b).

Clinical EFT is generally considered safe, with few accounts of abreactions or emotional flooding (Schulz, 2009; Flint, Lammers, & Mitnick, 2005). These benefits extend even to the most highly traumatized populations, such as combat veterans with PTSD (Church, Hawk, et al., 2013), Rwandan genocide orphans (Stone, Leyden, & Fellows, 2009), and Haitian earthquake survivors (Gurret Caufour, Palmer-Hoffman, & Church, 2012). In a recent critical survey of therapists conducted through Listservs such as Acceptance and Commitment Therapy, the Society for the Science of Clinical Psychology, and the Association of Behavioral and Cognitive Therapies regarding their use of EFT and similar methods, 42% reported employing these techniques (Gaudiano, Brown, & Miller, 2012).

Acupoint tapping has been evaluated in several studies using electroencephalogram (EEG) imaging to record brain waves. Swingle, Pulos, and Swingle (2004) used the EEG to compare the brain-wave frequencies of auto accident victims before and after they performed EFT, and noted a reduction of those frequencies associated with PTSD. Lambrou, Pratt, and Chevalier (2003) used acupressure tapping with claustrophobics, comparing them with a non-claustrophobic group, and found an increase after treatment in theta EEG frequencies, which are associated with relaxation, and a reduction in anxiety symptoms, which persisted on follow-up. Swingle (2010) found EFT to be beneficial in the treatment of seizure disorders. Regulation of the autonomic nervous system and the fear response has been found in fMRI studies using

acupuncture (Hui et al., 2005; Napadow et al., 2007; Fang et al., 2009), and similar effects for acupoint tapping have been proposed (Feinstein, 2010).

As EFT uses elements of efficacious therapies, some critics claim that any beneficial effects might be due solely to the merits of these therapies and that acupoint tapping makes no contribution to its success (Bakker, 2014). This is the so-called “purple hat” fallacy, which holds that a practitioner might employ a therapeutic method of demonstrated efficacy while wearing a purple hat and then attribute the success of treatment to the hat (Rosen & Davison, 2003). For this reason, dismantling studies that separate acupoint stimulation from the cognitive and exposure portions of the protocol have been recommended (Baker, Carrington, & Putilin, 2009), the first of which was performed by Fox (2013). In a randomized controlled trial with university students, the investigator administered the exposure portion of EFT to a control group, though without the cognitive “Setup Statement” that is a component of EFT. In an active control group, mindful breathing was substituted for tapping. The study found that the tapping group showed significantly better results on most measures.

A second dismantling study examined 126 schoolteachers at risk for burnout. EFT was compared to a control condition that included tapping on a sham location on the body coupled with the exposure component of EFT (Reynolds, 2015). Participants were assessed using the Maslach Burnout Inventory (MBI; Maslach, Jackson, & Leiter, 1996), which employs three scales: Emotional Exhaustion, Depersonalization, and Personal Accomplishment. The study reduced the possibility of cross-contamination between the two conditions by not randomizing participants within a single population. Instead, to minimize contact between experimental and control participants, the two samples were taken from different school districts with similar demographic profiles in the same county.

To ensure that the cognitive and exposure portions of both protocols were similar and thus isolate tapping as a viable and measurable technique, Reynolds (2015) gave both the EFT and control groups identical lists of troubling situations and cognitions that might contribute to burnout, and they were instructed to focus on them mentally while utilizing the procedure. Instead of using the Setup Statement and prescribed acupressure points of EFT, members of the control group were instructed to tap with an open right hand on the forearm of their left hand. The position of the hand doing the tapping is important, as there are acupoints at the tips of each finger. Thus this study controlled for the stimulation of those points by tapping with the undersides of the fingers instead of the fingertips, in contrast to an earlier study that unintentionally used fingertip points (Waite & Holder, 2003). The results of the Reynolds (2015) study showed that on all three indicators of burnout measured, EFT was superior to the sham tapping employed in the control group.

Waite and Holder (2003) compared three tapping conditions (EFT points, sham points, and points on a doll) to a non-tapping condition and found that participants in all three tapping groups showed significant improvements, while the non-tapping group did not. From this attempt at a dismantling study, the investigators concluded that EFT owed its efficacy to distraction and desensitization. The same findings have been reinterpreted as supporting EFT, however, as the investigators inadvertently engaged fingertip tapping points in the three groups that improved significantly (Baker, Carrington & Putilin, 2009; Pasahow, 2010). Another flaw in Waite and Holder's (2003) study was that it failed to adhere to the manualized form of the method and instead introduced novel variants to the protocol. Its departure from established research norms and its ambiguous results make this study an outlier in EFT research literature.

Another dismantling study performed by Church and Nelms (2015) recruited 34 participants with clinically verified “frozen shoulder” consisting of limited range of motion (ROM) measured in five different planes of arm movement. Participants were randomized into a wait list or one of two treatment groups. ROM, pain, and psychological conditions such as anxiety and depression were assessed before and after a 30-minute treatment session, and 30 days later. One treatment group received Clinical EFT, while the other received an identical cognitive and exposure protocol but with diaphragmatic breathing (DB) substituted for acupuncture. While the DB group showed improvements in pain and psychological distress after treatment, their results were not as pronounced as those of the tapping group. On follow-up the participants in the tapping group maintained their gains while those in the DB group did not. As in the Fox (2013) and Reynolds (2015) studies, these results indicate that acupoint tapping is an active ingredient in the therapeutic results obtained from EFT, and that its efficacy is not solely due to protocols such as exposure and cognitive restructuring that are shared with other therapies.

The fifth dismantling study used a convenience sample of 56 university students randomized into an EFT group and a control group that employed an identical protocol but with sham tapping points (Rogers & Sears, 2015). A stress test was administered before and after a single tapping session. The group that had tapped on actual acupuncture points exhibited a significantly greater reduction in stress ($p < .0001$). The study had several methodological weaknesses, however, such as the lack of a reliable and valid assessment and the administration of the intervention by one of the investigators.

Several meta-analyses of EFT have been published. The earliest was by Gilomen and Lee (2015) and examined studies published through 2012. It identified a moderate treatment effect,

but because its time frame did not include the dismantling studies summarized above, it speculated that treatment effects might be due to the non-specific benefits of any therapy. A later meta-analysis of EFT for the treatment of PTSD found very large treatment effects across seven RCTs (Sebastian & Papworth, 2015). It also found equivalence to established therapies and superiority of EFT to treatment as usual. A meta-analysis by Clond (2016) that did include the dismantling studies found a large effect size in the remission of anxiety symptoms after EFT treatment. This and other evidence points to EFTs clinically significant effects on a wide variety of psychological symptoms.

Depression as a Clinical Outcome

The National Survey on Drug Use and Health (NSDUH) defines a major depressive episode as: “A period of two weeks or longer during which there is either depressed mood or loss of interest or pleasure, and at least four other symptoms that reflect a change in functioning, such as problems with sleep, eating, energy, concentration, and self-image.” At least one depressive episode is reported by 15.7 million US adults annually, or 6.7% of the population aged 18 or older (Center for Behavioral Health Statistics and Quality, 2015).

Depression as a chronic condition is not simply a psychological problem; it has measurable physiological correlates, at the level of organic systems as well as at cellular and molecular levels. When compared with the brain scans of normal subjects, the scans of depression sufferers show much faster age-related loss of brain volume, as well as shrinkage of structures in the limbic system (Sheline, 2003). Telomeres, the molecular switches at the ends of chromosomes, have recently emerged as a primary marker of biological (as opposed to chronological) age. Shrinking telomeres indicate accelerated cellular aging. For this reason,

investigators parse the discrepancies between the biological and chronological ages of groups of subjects.

Recent studies examined the length of telomeres in depressed subjects to determine the association between depression and accelerated cellular aging, and found significant telomere loss (Hartmann, et al., 2013; Wikgren, 2011; Wolkowitz, Epel, Reus, & Mellon, 2010). The mean age of PTSD-positive participants in one of these studies (O'Donovan, 2011) was 30 years and, by that age, the telomeres of those who were depressed indicated a biological age 4.5 years older than non-depressed subjects. Telomere shortening is also associated with increased inflammation, premature mortality, and a complex of age-related illness including stroke, dementia, cardiac events, and diabetes (O'Donovan et al., 2011; Wolkowitz et al., 2010).

Study Aim

We sought to collect information pertaining to EFT methods most recently used in clinical trials, both randomized and observational, and to examine their relative efficacy in reducing depression in different samples. During the preliminary search of the literature, we organized a review of the studies published in the last 10 years. We chose this time frame as an appropriate span for forming an accurate picture of the state of the research evidence. Further, we postulated that a sufficient number of studies would have been published during that time to allow for meta-analytic calculations. As an easily-taught method that is used both as a self-help technique and in clinical practice, we proposed to identify whether or not there is an empirical basis for EFTs widespread use for depression (Church, 2013b).

Methods

Study Selection

The overall objective of study selection was to collect reports published in peer-reviewed journals that examined depression symptom levels before and after EFT treatment both in clinical and non-clinical populations. A literature search for English-language papers was performed using MEDLINE/PubMed, PsycINFO, Google Scholar, and references from the retrieved papers. The search is current through June of 2015. Keyword searches included “Emotional Freedom Technique” and “depression,” “Emotional Freedom Techniques” and “depression,” and “EFT” and “depression.” We also included “post-traumatic stress disorder” and “PTSD” as search terms, as we noted that studies of the effect of EFT on PTSD often include an examination of psychological symptoms (including depression) through the use of a standardized questionnaire. Professional organizations in the field, and individual authors, were queried in order to obtain information about unpublished “file drawer” studies as well as studies “in press.”

Though some researchers (e.g., Cochrane Collaborative) view the RCT as the only acceptable evidence for treatment outcome, many systematic reviews are indeterminate because they include insufficient numbers of RCTs whilst rejecting large numbers of non-randomized controlled studies. We decided to include all studies published and relevant to our aim(s), independent of their research design, in order to increase the sample size and provide clinically informative results.

Of RCTs, only those that assessed depression as an outcome measure and met the APA Division 12 criteria were included. To meet the Division 12 efficacy criteria as a “well-established treatment,” two independent research teams must have demonstrated in a peer-

reviewed report that the treatment is either (a) more effective than a placebo or (b) that it is equivalent to another treatment whose efficacy has been established (Chambless & Hollon, 1998; Chambless et al., 1998). In order to be designated and “efficacious” method, the treatment must also be standardized (which is usually taken to mean that the practitioner delivering the treatment adhered to manualized procedures) and the client population must be clearly specified so the study’s generalizability can be inferred. To achieve a “probably efficacious treatment” designation, only one study meeting the above criteria is required. The “probably efficacious” criteria can be met by two studies in which the control condition is a wait-list group instead of a placebo condition or an established treatment. The two studies may be by the same investigator or different investigators. The Division 12 criteria were therefore used as a convenient published standard allowing us to screen for the quality of RCTs.

Study Eligibility

The first author (JN) screened the abstracts of all publications obtained by the search strategy. Studies meeting the following inclusion criteria were selected for the meta-analysis: (a) at least one session of EFT, (b) reporting of interval or ratio data, (c) use of valid and reliable psychometric questionnaires; (d) pre and post-treatment depression level data presented, and (e) sufficient reporting of study results (e.g., means and standard deviations) to allow for effect size computation. Some studies utilized both repeated measures designs (before and after EFT), as well as comparisons (EFT versus active treatment or wait list). The absence of a control group was not an exclusion criterion because calculation of effect size can be performed on pre-post changes. However, given that between-group and within-group analyses are methodologically different (Shrier, 2007), two separate analyses were conducted. In their review, Shrier et al. (2007) argue for the inclusion of non-RCTs in meta-analyses. They conclude that the benefits

outweigh the disadvantages and point out that the effect sizes found in outcome studies are generally similar to those in RCTs. We did not require an independent observer-rated diagnosis of depression since we believe that self-reported depressive symptoms are more clinically significant to depression sufferers than observer-rated measures, and the discrepancies between the two are anyway usually minor (Enns, Larsen, & Cox, 2000).

One study by Nemiro & Papworth (2015) assessed general mental health, with depression and anxiety measured in sub-scales. The investigators only provided the general mental health scale, and despite their depression results being requested several times for this meta-analysis, they did not provide this data and we were therefore unable to include it. We also excluded several studies which re-analyzed data from studies already included. Excluded studies were Church (2013) which was a detailed analysis of the interaction of pain, anxiety, and depression using data extracted from Church, Hawk, et al. (2013); Hartung and Stein (2012) which compared office to telephone sessions, and Stein and Brooks (2011) which compared EFT delivered by life coaches versus licensed mental health professionals.

Data Coding

For all papers selected, the full articles were obtained and inspected to assess their relevance based on the preplanned criteria for inclusion. Data were extracted using a predesigned data collection form: (1) number of subjects, (2) geographic origin of the study, (3) treatment delivery format, (4) description of population, (5) mean age, (6) percentage of women, (7) assessment measures, (8) homework, (9) protocol length, (10) trial context, and (11) summary statistics required for computation of effect sizes. Data were then reviewed by the first and second authors (JN and LC) to reach an agreement. Any disagreements were resolved by an outside reviewer.

Besides computing the total average effect size of EFT intervention for depression, we also calculated effect sizes across multiple time points. We did this by structuring an estimating equation with weights assigned to each study based on the number of participants used in the analysis. Time points analyzed included posttest, less than 90 days, and 90 days or greater. Ambiguity concerning study sample sizes (i.e., unspecified attrition) was solved by the conservative approach of using the smallest number for which there was clear documentation.

Depression was the dependent variable. From the studies selected for the meta-analysis, we extracted all available psychometric data, which were available from a variety of questionnaires: Symptom Assessment-45 (SA-45; Davison et al., 1997), Beck Depression Inventory II (BDI II; Beck, Steer, & Carbin, 1988), Depression Anxiety Stress Scales (DASS; Henry & Crawford, 1995), Hospital Anxiety and Depression Scale (HADS; Zigmond & Snaith, 1983), and the Profile of Mood States (POMS; Curran, Andrykowski, & Studts, 1995) depression subscale.

Statistical methods

To assess clinically relevant treatment effects, the percentage change in depressive symptoms before and after treatment was calculated using normed T-scores or whatever other scoring method was specified in the relevant assessments. Effect sizes were measured using Cohen's *d*. When the necessary data were available, all effect sizes were calculated directly using the following formula: $d = (M1 - M2)/S$, where *M1* is the pretest mean, *M2* the posttest or follow-up mean, and *S* the standard deviation for the pooled sample. We used Cohen's *d* rather than Hedge's *g* because it is more commonly used in meta-analyses and therefore allowed us to compare results with meta-analyses of other treatments such as psychopharmacology and psychotherapy. For RCTs, the effect size was first calculated only for the EFT treatment group.

For within-group outcome studies without a control group, baseline scores instead of control group scores were used in the formula. If these data were not provided, d was estimated using conversion equations for significance tests. If an RCT had an active control, such as EMDR, a comparison was then made between the treatment effectiveness of the two therapies. The overall mean effect size for all of the studies combined was weighted by the variance of the studies, considering both standard deviations and number of subjects. Effect sizes are usually interpreted using Cohen's original convention of small (<0.2), medium (0.5), and large (>0.8) treatment effects.

Results

We identified 20 studies for inclusion in the meta-analysis. Eight studies investigated EFT outcomes only; the other 12 were RCTs. The pooled sample for outcome studies comprised 461 subjects, while the pooled RCT sample consisted of 398 participants who received EFT treatment. Study characteristics are listed in Table 1. Tables 2 and 3 outline the reported results for outcome studies and RCTs respectively. Several types of psychological surveys were employed to measure depressive symptoms in the studies included in this meta-analysis.

We first calculated effect sizes as Cohen's d comparing depression scores at time points after EFT intervention with pre-intervention scores. We then stratified time points into posttest, <90 days, and 90 days or more (Tables 4, 5 and 6). We calculated weighted mean effect size for each stratum. For the multivariate model, we took as the one comparison per study the 90-day assessment in an EFT group if the study extended 90 days or longer. The outcome variable was depression scale scores. For the studies included in the meta-analysis multivariate model, we calculated as the outcome the standardized mean difference in 90-day follow-up versus pre-intervention scores.

Tables 2 and 3 illustrate the effect sizes pretest and posttest for all studies. Tables 4, 5 and 6 illustrate time point-specific measurements. The letter d represents Cohen's d , while $SE(d)$ represents the standard errors in each respective Cohen's d measurement. Finally, $p(d)$ represents the probability of observing this or more larger effect size by chance. As can be seen from Tables 2 and 3, the majority of studies demonstrated a statistically significant change in depression compared to pretest. The weighted effect sizes for outcome studies is 0.7046. The pretest-posttest effect size for RCTs is 1.8498. Overall, the pre-post effect size is 1.3087.

The same pattern holds for measurements taken at less than 90 days (Table 5) where five of the eight studies reported a statistically significant change. Table 6 reports measurements at 90 days and greater. All 90-day or less versus pretest effect sizes were reported as being at least 0.29, with effects ranging from 0.29 to 3.78. For less than 90 days follow up versus pretest comparisons, the outcome studies have an effect size of 0.8102. The RCTs effect size is 1.3832. The weighted effect size for these eight studies is 1.2119. Similarly, the effect size for 90 days follow up or greater versus pretest is 0.5253 and 1.73 for outcome and RCT studies, respectively.

We also assessed heterogeneity using a Q statistic to determine which type of model should be used for computing estimates of effect sizes 90 days or more after treatment. We found a Q of 22.39 ($p = 0.0004$), which indicates significant heterogeneity among samples within this time period. We also calculated the between-study variance (τ^2) as 0.4505. Based on this calculation, we used a random effects model to estimate the effect size due to EFT as $d = 1.11$ ($p = 0.0007$). This effect size is very large and indicates a consistent depression score difference from baseline to after EFT treatment, and that the effects of EFT treatment are lasting, up to and through 90 days after the intervention. The weighted mean of reductions in depressive symptoms was measured and found to be -41%.

Six RCTs used an active treatment (not wait list) as the comparison group. Two provided treatment as usual (TAU), one provided supportive interviews (SI), one provided diaphragmatic breathing (DB), one provided Eye Movement Desensitization and Reprocessing (EDMR) therapy, and one provided Sleep Hygiene Education (SHE). Table 7 illustrates the statistical differences between EFT and DB, SI, EDMR, and SHE. We did not include treatment as usual (TAU) as an active control group. EFT was more efficacious than DB and SI in the posttest measurements ($p = 0.06$ vs. DB; $p < 0.001$ vs. SI). EFT was more efficacious than SHE at the 9th week assessment ($p = 0.036$). We found no statistically significant difference between EFT and EDMR at posttest. Most studies reported that no adverse reactions were noted, and that participant distress decreased rapidly across all symptom domains simultaneously.

Discussion

This study evaluated the impact that EFT has on depression in general and among population subgroups. We found that EFT was efficacious in mitigating depression in a variety of demographic groups. A meta-analysis of psychopharmacology for mild depression found Cohen's $d = .20$ (Fournier et al., 2010), while high-quality trials of psychotherapy found an effect size of $d = .22$ (Cuijpers, van Straten, Bohlmeijer, Hollon, & Andersson, 2010). The effect size found for EFT in this meta-analysis exceeded the treatment effect typically found in psychopharmacology and psychotherapy trials. Further, we discovered that the impact of EFT, especially in interventions performed over multiple sessions, endures up to and surpasses 90 days, even up to 6 months post-intervention.

The findings of two studies that were excluded since they were re-analyses of other work are nonetheless of clinical interest. Hartung and Stein (2012) performed a reanalysis of Church, Hawk, et al., (2013) comparing office to telephone phone sessions; it found that in-person EFT

treatment was significantly more effective. Stein and Brooks (2011) conducted a third reanalysis of Church, Hawk, et al. (2013) to determine if there is a difference between EFT delivered by life coaches versus licensed mental health professionals. This study found that for depression the difference was not statistically significant. This indicates that EFT is as effective when used by life coaches as by mental health professionals, pointing to the method's utility in settings with limited access to highly trained clinicians.

Though the number of studies in which EFT was compared to established treatments in this meta-analysis was small, in all cases EFT was found to have equal or greater efficacy. EFT was also more efficacious than treatment as usual or the passage of time. One of the studies included in this analysis (Church & Nelms, 2016) was a dismantling study. It was carefully designed to determine whether or not the acupoint tapping described in EFT was an active ingredient in the treatment protocol. It compared a tapping group to a group that performed an identical protocol but with diaphragmatic breathing (DB) substituted for tapping. While positive effects for the DB group were found, they were not as great as for the tapping group, and they did not persist over time. An earlier meta-analysis found that EFT produced a moderately large treatment effect, but based on the limited number of studies then available to the authors, it was impossible to determine whether treatment effects were due to tapping, or to the non-specific factors found in any therapy (Gilomen & Lee, 2014). The current study allowed us to test this hypothesis, and on the basis of the included study, as well as several other dismantling studies published subsequent to Gilomen and Lee (2014), it appears that tapping is an active ingredient in EFT and not an inert placebo.

As did the extensive review by Shrier et al. (2007) we found that there is not a marked difference in effect sizes between outcome studies and RCTs provided that sufficient numbers of

both are available for analysis. We therefore suggest that including outcome studies in meta-analyses enriches their utility for clinical decision-making. EFTs clinical utility and generalizability is supported by several characteristics: The brief treatment time frames found in the studies examined, ranging from one to 10 sessions; its efficacy in group format as well as individual counseling sessions; the absence of abreactions in participants; the unremarkable dropout rate, and the effects across demographic groups, countries, and treatment settings.

The meta-analysis was strengthened by several factors. One the use of the same assessment, the SA-45, in 13 of the 20 studies included, enabling conclusions to be drawn from a variety of demographic samples. Another was the stated use of *The EFT Manual* (Craig & Fowlie, 1995; Church, 2013) in most of the reports, indicating consistent application of the therapeutic method. Reported studies were published in a variety of journals, were conducted by a variety of investigators, assessed demographically diverse samples including participants in several different countries, and used assessments other than the SA-45 such as the HADS and BDI. These factors all argue for the generalizability of the results, subject to the limitations outlined below.

Limitations

There are several limitations to this study and virtually any other meta-analysis. Findings from this meta-analysis must be interpreted with caution given limitations of meta-analysis in general and of data collected for this analysis in particular. A critical issue for this meta-analysis, as is true of any systematic review, was deciding which trials or studies to include and which to exclude. As noted, we included all studies published and relevant to our aim(s), independent of their research design, in order to increase the number of studies and participants. All outcome studies including the ones analyzed in this paper have the limitation that the improvements they

note may be the nonspecific results of any treatment.

As in any review of studies in a given area, it is possible that studies with non-significant results are underreported. Publication bias, the possibility that only studies with significant outcomes have been published and those with null outcomes have not, skew the results of meta-analyses. While we did contact the professional organizations in the field in an attempt to locate unpublished studies, it is possible that some exist unknown to us or to those organizations. It is further important to note that, for some studies, analysis was based on relatively few subjects.

We searched studies in the most important databases for psychology (PsychInfo) and medicine (Medline). Other databases (e.g. CINAHL) were not screened, however, and this may be a limitation to the generalizability of our results, given that other studies might be available in these databases. A high degree of heterogeneity was found; factors that might have contributed to this were the large number of studies, the diversity of populations from clinic clients with major depressive disorder to veterans with PTSD to non-clinical workshop participants, and the varied treatment time frames ranging from one to 10 sessions.

Another major limitation generally present in studies, including those of EFT, is a substantial loss to follow-up. Many studies in our analysis experienced up to 50% loss to follow-up after the intervention period. Loss to follow-up occurs for a variety of reasons including: lack of compensation for participation, length of time required to complete follow-up assessments, and the difficulty that participants experience in keeping appointments among life's other demands. Regardless of the reason, this factor may introduce bias into the measurement of effect or tests of significance. It is impossible estimate the effect of loss to follow-up on the studies analyzed; therefore, we cannot be certain as to the difference complete follow-up would have made in the treatment or control groups. However, the dropout rate between the start and end of

the EFT intervention period is consistent with the rate of about 20% found in psychotherapy trials (Ekers, Richards, & Gilbody, 2008).

Our meta-analysis also included relatively few studies comparing EFT to another established treatment. Most studies used a wait list or treatment as usual group as a control. Further studies are needed to gauge the efficacy of EFT relative to established methods. Also, many of the studies used depression as one of several outcomes, including anxiety and PTSD. Most used participant self-report, rather than observer-rated measures. Only two used a categorical diagnosis of major depressive disorder. These limitations make it difficult to generalize the findings of this study to populations that typically seek treatment for severe depression. Instead, only the more conservative conclusion that EFT reduces depressive symptoms in a wide sample of demographic groups can be drawn from this analysis.

Though EFT is generally standardized and manualized throughout the world, the length and number of EFT sessions offered to study participants varies. We calculated the effect of an EFT intervention across studies; however, we were unable to determine if the effect size differed across intervention lengths as study lengths varied greatly across the group of studies. Two approaches could be taken to enhance the ability of researchers to measure the precise effect of EFT over various time points. First, EFT intervention could be further normalized to establish a standard length and time for all interventions. A further normalization of EFT time points would allow for the assessment of EFT effectiveness across virtually any health outcome.

Secondly, daily or weekly monitoring could be performed to assess the change in a health outcome at normalized intervals (e.g. posttest, 3 days, 1 week, 2 weeks, 30 days, 60 days, 90 days, etc.). Such an approach would allow subjects involved in an EFT study to have their progress monitored in any health outcome more frequently, thus providing more data points for

study. With more data points available for each participant, even those who drop out could at least present one data point, though they failed to complete the entire intervention.

Furthermore, if many data points are taken across the study period, interpolation, imputing, or other modeling techniques could be employed to estimate the effect at any time point and for any participant that may be missing. Based on existing data, this type of modeling and smoothing is not feasible.

Conclusions

The current study used meta-analytical methods to examine the efficacy of EFT for depression. It identified a large treatment effect across a wide demographic spectrum, with participants maintaining their gains over time. EFT was effective in brief treatment time frames ranging from one to 10 sessions, and when delivered in group treatment format as well as individual therapy sessions. It found greater efficacy for EFT than typically found in trials of psychopharmacology and psychotherapy. EFT was more efficacious than usual care and than most active controls, and was as effective as EMDR, another evidence-based treatment, in one study. As a safe and reliable self-help method, EFT demonstrates clinical utility as a low-cost non-drug treatment for depression in a wide variety of settings and demographic groups. Based on the RCTs in our meta-analysis, EFT unequivocally meets the APA Division 12 criteria as an efficacious therapy for depression.

Competing interests: The authors declare that they have no competing interests.

Acknowledgments: The research was supported by donations to the National Institute for Integrative Healthcare (niih.org), which especial thanks Nick Ortner, and Robert and Lynne Hoss, for their contributions.

References

Baker, B. Hoffman, C. (2014). Emotional Freedom Techniques (EFT) to reduce the side effects associated with tamoxifen and aromatase inhibitor use in women with breast cancer: A service evaluation. *European Journal of Integrative Medicine*

Brattberg, G. (2008). Self-administered EFT (Emotional Freedom Techniques) in Individuals With Fibromyalgia: A Randomized Trial. *Integrative Medicine*, 7 (4): 30-35.

Church, D., Brooks, A. (2010). The Effect of a Brief Emotional Freedom Techniques Self-Intervention on Anxiety, Depression, Pain, and Cravings in HealthCare Works. *Integrative Medicine*, 9(5): 40-43.

Church, D., Brooks, A., (2013). The Effect of EFT (Emotional Freedom Techniques) on Psychological Symptoms in Addiction Treatment: A Pilot Study. *Journal of Scientific Research & Reports*, 2(1): 315-213. Article no. JSRR.2013.021.

Church, D. (2010). The treatment of combat trauma in veterans using EFT: A pilot protocol. *Traumatology*, 15(1), 45-55.

Church, D., Geronilla, L., Dinter, I. (2009). Psychological Symptom Change in Veterans After Six Sessions of Emotional Freedom Techniques (EFT): An Observational Study. *Wholistic Health Publications*, 9 (1). 14 pages.

Church, D., & Nelms, J. (2015). Psychological change in a population with frozen shoulder: A randomized controlled dismantling study of clinical EFT (Emotional Freedom Techniques). Presented at the ACEP conference, May 31, 2015.

Church, D., De Asis, M.A., Brooks, A.J. (2012). Brief Group Intervention Using Emotional Freedom Techniques for Depression in College Students: A Randomized Controlled Trial. *Depression, Research, and Treatment*, 2012, 1-7. Article ID: 257172.

Church, D., Hawk, C., Brooks, A.J., Toukolehto, O., Wren, M., Dinter, I., Stein, P. (2013). Psychological Trauma Symptom Improvement in Veterans Using Emotional Freedom Techniques – A Randomized Controlled Trial. *J Nerv Ment Dis*, 201: 153-160.

Church, D., Sparks, J.D., Clond, M. (2014). EFT (Emotional Freedom Techniques) and Resilience in Veterans at Risk for PTSD: A Randomized Controlled Trial. Elsevier Editorial System™ for EXPLORE: *The Journal of Science and Healing*.

Church, D., Yount, G., Brooks, A. J. (2012). The Effect of Emotional Freedom Techniques on Stress Biochemistry – A Randomized Controlled Trial. *J Nerv Ment Dis*, 200: 891-896).

Church, D., Yount, G., Rachlin, K., Fox, L., & Nelms, J. (2015). Epigenetic effects of PTSD remediation in veterans using Clinical EFT (Emotional Freedom Techniques): A randomized controlled trial. Data presented at the conference of the Association for Comprehensive Energy Psychology, Virginia Beach, VA, May 30, 2015.

Cuijpers, P., van Straten, A., Bohlmeijer, E., Hollon, S. D., & Andersson, G. (2010). The effects of psychotherapy for adult depression are overestimated: a meta-analysis of study quality and effect size. *Psychological Medicine*, 40(02), 211-223.

Ekers, D., Richards, D., & Gilbody, S. (2008). A meta-analysis of randomized trials of behavioural treatment of depression. *Psychological Medicine*, 38(05), 611-623.

Enns, M. W., Larsen, D. K., & Cox, B. J. (2000). Discrepancies between self and observer ratings of depression: the relationship to demographic, clinical and personality variables. *Journal of affective disorders*, 60(1), 33-41.

Geronilla, L., McWilliams, M., & Clond, M. (2014). EFT (Emotional Freedom Techniques) remediates PTSD and psychological symptoms in veterans: A randomized controlled replication trial. Presented at the Grand Rounds, Fort Hood, Killeen, Texas, April 17.

Fournier, J. C., DeRubeis, R. J., Hollon, S. D., Dimidjian, S., Amsterdam, J. D., Shelton, R. C., & Fawcett, J. (2010). Antidepressant drug effects and depression severity: a patient-level meta-analysis. *JAMA*, *303*(1), 47-53.

Hartmann, N., Boehner, M., Groenen, F., & Kalb, R. (2010). Telomere length of patients with major depression is shortened but independent from therapy and severity of the disease. *Depression and Anxiety*, *27*(12), 1111-1116.

Lee, J.H., Chung, S.Y. Kim, W.O. (2015). A Comparison of Emotional Freedom Techniques – Insomnia (EFT-I) and Sleep Hygiene Education (SHE) for Insomnia in a Geriatric Population: A Randomized Controlled Trial. *Energy Psychology: Theory, Research, and Treatment*, *7*(2), 22-29.

Karatzias, T., Power, K., Brown, K., McGoldrick, T., Begum, M., Young, J., Loughran, P., Adams, S. (2011). A Controlled Comparison of the Effectiveness and Efficiency of Two Psychological Therapies for Posttraumatic Stress Disorder: Eye Movement Desensitization and Reprocessing vs. Emotional Freedom Techniques. *J Nerv Ment Dis*, *199*: 372-378

Nemiro, A., & Papworth, S. (2015). Efficacy of two evidence-based therapies, Emotional Freedom Techniques (EFT) and Cognitive Behavioral Therapy (CBT) for the treatment of gender violence in the Congo: A randomized controlled trial. *Energy Psychology: Theory, Research, and Treatment*, *7*(2), 13–25. doi:10.9769/EPJ.2015.11.1.AN

Rowe, J.E. (2005). The Effects of EFT on Long-Term Psychological Symptoms. *Counseling and Clinical Psychology Journal*, (3): 104-111.

Shrier, I., Boivin, J. F., Steele, R. J., Platt, R. W., Furlan, A., Kakuma, R., ... & Rossignol, M. (2007). Should meta-analyses of interventions include observational studies in addition to

randomized controlled trials? A critical examination of underlying principles. *American Journal of Epidemiology*, 166(10), 1203-1209.

Stapleton, P., Church, D., Sheldon, T., Porter, B., & Carlopio, C. (2013). Depression symptoms improve after successful weight loss with emotional freedom techniques. *ISRN Psychiatry*, 2013, 7.

Stapleton, P., Devine, S., Chatwin, H., Porter, B., Sheldon, T. (2014) A Feasibility Study: Emotional Freedom Techniques For Depression In Australian Adults. *Current Research in Psychology*, 5(1): 19-33.

Stewart, A., Boath, E., Carryer, A., Walton, I., Hill, L. (2013) Can Emotional Freedom Techniques (EFT) be effective in the treatment of emotional conditions? Results of a service evaluation in Sandwell. *Journal of Psychological Therapies in Primary Care*, 2, 71-84.

Wolkowitz, O. M., Epel, E. S., Reus, V. I., & Mellon, S. H. (2010). Depression gets old fast: do stress and depression accelerate cell aging?. *Depression and Anxiety*, 27(4), 327-338.

Table 1. Study Characteristics

Number	Authors	Mean Age	% Women	Country	Treatment	Population	Instrument	Delivery Method
1	Baker & Hoffman, 2015	55	100	UK	EFT Shortcut Basic Recipe	Women with breast cancer taking hormones Workshop participants with addiction issues	POMS	Group
2	Church & Brooks, 2013	54	72	US	Clinical EFT	Health care personnel	SA-45	Group
3	Church & Brooks, 2010	48	76	US	Clinical EFT	Veterans	SA-45	Group
4	Church, Geronilla, Dinter, 2009	NR	43	US	Clinical EFT	Convenience sample of workshop participants	SA-45	Group
5	Rowe, 2005	53	74	US	Clinical EFT	Major Depressive Disorder clients at a university clinic	SA-45	Group
6	Stapleton, Devine, Chatwin, Porter, Sheldon, 2014	NR	82	Australia	Clinical EFT	National Health Service hospital patients	BDI II; DASS	Group
7	Stewart, Boath, Carryer, Walton, Hill, 2013	45	77	UK	Clinical EFT	Veterans and family with PTSD	HADS	Group
8	Church, 2010	NR	36	US	Clinical EFT	Fibromyalgia patients on sick leave	SA-45	Individual
9	Brattberg, 2008	44	100	Sweden	Clinical EFT	Veterans with PTSD	HADS	Group
10	Church, Hawk, Brooks, Toukolehto, Wren, Dinter, Stein, 2013	52	10	US	Clinical EFT	Veterans at risk of PTSD	SA-45	Individual
11	Church, Sparks, Clond, 2014	56	44	US	Clinical EFT	Nonclinical subjects	SA-45	Individual
12	Church, Yount, Brooks, 2012	49	82	US	Clinical EFT	Patients with Major Depressive Disorder	SA-45	Individual
13	Stapleton, Sheldon, Porter, Whitty, 2011	NR	89	Australia	Clinical EFT		SA-45	Group

14	Karatzias et al., 2011	40	52	UK	EFT Movie Technique	National Health Service psychotherapy referrals	HADS	Individual
15	Lee, Chung, Kim, 2015	80	100	South Korea	Clinical EFT	Senior insomnia patients	GDS-K	Group
16	Church, De Asis, Brooks, 2012	17	80	Philippine s	Clinical EFT	Psychology students	BDI	Group
17	Geronilla, McWilliams, Clond, 2014	50	19	US	Clinical EFT	Veterans with PTSD	SA-45	Individual
18	Church & Nelms, 2015	54	63	US	Clinical EFT	Adults with frozen shoulder	SA-45	Individual
19	Church, Yount, Rachlin, Fox, Nelms, 2015	60	31	US	Clinical EFT	Veterans	SA-45	Individual
20	Stapleton, Church, Sheldon, Porter, Carlopio, 2013	NR	89	Australia	Clinical EFT	Overweight and obese adults	SA-45	Group

Note. BDI II = Beck Depression Inventory II; DASS = DASS Depression Anxiety Stress Scales; DB = diaphragmatic breathing; EFT = Emotional Freedom Techniques; EMDR = Eye Movement Desensitization and Reprocessing; GDS-K = Geriatric Depression Scale in Korea; HADS = Hospital Anxiety and Depression Scale; NT = no treatment; POMS = Profile of Mood States; SA-45 = Symptom Assessment-45; SHE = sleep hygiene education; SI = supportive interview; TAU = treatment as usual; WL = wait list; NR = not reported

Table 2. Pretest and Posttest Percent Change and p-values for Outcome Studies

Number	Authors	Analytical Sample Size	Country	Depression Measurement (Instrument)	Pre vs Post % Change	p-value
1	Baker & Hoffman, 2015	41	UK	POMS	-16%	0.006
2	Church & Brooks, 2013	39	US	SA-45	-26%	0.024
3	Church & Brooks, 2010	216	US	SA-45	-60%	0.001
4	Church, Geronilla, Dinter, 2009	7	US	SA-45	-34%	0.001
5	Rowe, 2005	102	US	SA-45	-60%	0.001

6	Stapleton, Devine, Chatwin, Porter, Sheldon, 2014	6	Australia	BDI II	-29%	0.028
7	Stewart, Boath, Carrier, Walton, Hill, 2013	39	UK	HADS	-48%	0.001
8	Church, 2010	11	US	SA-45	-87%	0.005

Note. BDI II = Beck Depression Inventory II; DASS = DASS Depression Anxiety Stress Scales; DB = diaphragmatic breathing; EFT = Emotional Freedom Techniques; EMDR = Eye Movement Desensitization and Reprocessing; GDS-K = Geriatric Depression Scale in Korea; HADS = Hospital Anxiety and Depression Scale; NT = no treatment; POMS = Profile of Mood States; SA-45 = Symptom Assessment-45; SHE = sleep hygiene education; SI = supportive interview; TAU = treatment as usual; WL = wait list; NR = not reported

Table 3. Pretest and Posttest Percent Change in Symptoms and p-values After EFT in RCTs

Number	Authors	Analytical Sample Size	Country	Depression Measurement (Instrument)	Pre vs Post % Change	p-value
1	Brattberg, 2008	30	Sweden	HADS	-29%	0.02
2	Church, Hawk, Brooks, Toukolehto, Wren, Dinter, Stein, 2013	49	US	SA-45	-58%	0.001
3	Church, Sparks, Clond, 2014	18	US	SA-45	-47%	0.001
4	Church, Yount, Brooks, 2012	28	US	SA-45	-49%	0.001
5	Stapleton, Sheldon, Porter, Whitty 2011	96	Australia	SA-45	-23%	0.01

6	Karatzias et al., 2011	23	UK	HADS	-28%	0.001
7	Lee, Chung, Kim, 2015	10	South Korea	GDS-K	-60%	0.005
8	Church, De Asis, Brooks, 2012	9	Philippines	BDI	-74%	<0.05
9	Geronilla, McWilliams, Clond, 2014	58	US	SA-45	-48%	0.001
10	Church & Nelms, 2015	16	US	SA-45	-44%	0.001
11	Church, Yount, Rachlin, Fox, Nelms, 2015	16	US	SA-45	-38%	0.001
12	Stapleton, Church, Sheldon, Porter, Carlopio, 2013	45	Australia	SA-45	-21%	0.001

Table 4. Posttest Effect Sizes for Both Outcome Studies and RCTs

Study Number	Study	Type	d	SE(d)	p(d)
1	Baker & Hoffman, 2015	Outcome	0.19	0.29	0.52
2	Church & Brooks, 2013	Outcome	0.26	0.23	0.2
3	Church & Brooks, 2010	Outcome	0.75	0.1	<0.001
4	Church, Geronilla, Dinter, 2009	Outcome	1.1	0.57	0.06
5	Rowe, 2005	Outcome	\	\	\
6	Stapleton, Devine, Chatwin, Porter, Sheldon, 2014	Outcome	0.04	0.58	0.95
7	Stewart, Boath, Carryer, Walton, Hill, 2013	Outcome	0.84	0.31	0.01
8	Church, 2010	Outcome	2.64	0.57	<0.001

9	Brattberg, 2008	RCT	0.62	0.26	0.02
10	Church, Hawk, Brooks, Toukolehto, Wren, Dinter, Stein, 2013	RCT	8.02	0.78	<0.001
11	Church, Sparks, Clond, 2014	RCT	3.11	0.68	<0.001
12	Church, Yount, Brooks, 2012	RCT	1.12	0.29	<0.001
13	Stapleton, Sheldon, Porter, Whitty 2011	RCT	0.28	0.22	0.21
14	Karatzias et al., 2011	RCT	0.69	0.39	0.08
15	Lee, Chung, Kim, 2015	RCT	1.41	0.41	<0.001
16	Church, De Asis, Brooks, 2012	RCT	7.57	1.29	<0.001
17	Geronilla, McWilliams, Clond, 2014	RCT	1.93	0.3	<0.001
18	Church & Nelms, 2015	RCT	0.88	0.37	0.02
19	Church, Yount, Rachlin, Fox, Nelms, 2015	RCT	0.9	0.36	0.01
20	Stapleton, Church, Sheldon, Porter, Carlopio, 2013	RCT	0.37	0.23	<0.001

Weighted effect size—Outcome studies=0.7046

Weighted effect size—RCTs=1.8498

Weighted effect size—All studies=1.3087

Table 5. Effect Sizes Measured at Less Than 90 Days

Study Number	Study	Type	d	SE(d)	p(d)
1	Baker & Hoffman, 2015	Outcome	0.29	0.3	0.34
6	Stapleton, Devine, Chatwin, Porter, Sheldon, 2014	Outcome	0.79 2	0.6 0.69	0.2 0.01
7	Stewart, Boath, Carryer, Walton, Hill, 2013	Outcome	2.3	0.87	0.02
8	Church, 2010	Outcome	1.1	0.46	0.02
11	Church, Sparks, Clond, 2014	RCT	3.78	0.76	<0.01
15	Lee, Chung, Kim, 2015	RCT	0.94	0.38	0.02
18	Church & Nelms, 2015	RCT	1.1	0.38	<0.01
19	Church, Yount, Rachlin, Fox, Nelms, 2015	RCT	0.53	0.35	0.13

Weighted effect size—Outcome studies=0.8102

Weighted effect size—RCTs=1.3832

Weighted effect size—All studies=1.2119

Table 6. Effect Sizes for 90 Days and Greater

Study Number	Study	Type	d	SE(d)	p(d)
2	Church & Brooks, 2013	Outcome	0.32	0.25	0.2
4	Church, Geronilla, Dinter, 2009	Outcome	1.24	0.58	0.04
8	Church, 2010	Outcome	0.75	0.44	0.09
11	Church, Sparks, Clond, 2014	RCT	4.32	0.83	<0.01
19	Church, Yount, Rachlin, Fox, Nelms, 2015	RCT	0.87	0.36	0.02
20	Stapleton, Church, Sheldon, Porter, Carlopio, 2013	RCT	0.87	0.23	<0.01

Weighted effect size—Outcome studies=0.5253

Weighted effect size—RCTs=1.73

Weighted effect size—All studies=1.1093

Table 7. Significance Tests for Difference Between EFT and Other Active Treatments

Control Treatment	Follow-up	Study	<i>p</i> for difference
DB	Pre	Church & Nelms, 2015	0.16
	Post		0.06
	30-day		0.2108
SI	Post	Church, Yount, Brooks, 2012	<0.0001
EMDR	Post	Karatzias et al., 2011	0.724
SHE	9th week	Lee, Chung, Kim, 2015	0.036

Note. DB = diaphragmatic breathing; EMDR = Eye Movement Desensitization and Reprocessing; SHE = sleep hygiene education; SI = supportive interview

Accepted manuscript